



Correction to: Propofol but not dexmedetomidine produce locomotor sensitization via nitric oxide in rats

Tuğçe Uskur¹ · Ayşe Özçetin Şenöz² · Burcu Çevreli² · Aydın Barlas³ · Tayfun Uzbay^{2,4}

Published online: 20 November 2020

© Springer-Verlag GmbH Germany, part of Springer Nature 2020

Correction to: Psychopharmacology

<https://doi.org/10.1007/s00213-020-05707-5>

Page 3, Table 1, second line of table: The control group was injected with saline plus saline, not saline plus propofol. Thus injections in the second line of Table 1 should be corrected as saline saline.

The original article has been corrected.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at <https://doi.org/10.1007/s00213-020-05707-5>

✉ Tayfun Uzbay
tuzbay@uskudar.edu.tr; uzbayt@yahoo.com

¹ Faculty of Medicine, Department of Medical Pharmacology, Beykent University, Istanbul, Turkey

² Neuropsychopharmacology Application and Research Center (NPFUAM), Üsküdar University, Istanbul, Turkey

³ School of Medicine, Department of Pharmacology, Altınbaş University, İstanbul, Turkey

⁴ Faculty of Medicine, Department of Medical Pharmacology, Üsküdar University, İstanbul, Turkey